

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

62 COPEIA

A LIST OF PHILIPPINE FISHES.

During the past summer the collection of Philippine Fishes in the Commercial Museums of Philadelphia was kindly placed in my hands for examination and study by the Director, Dr. William P. Wilson. No localities were found with the specimens, and only a few native names, so they are admitted here as simply from the Philippine Islands. Species new to the fauna are indicated by the prefixed asterisk.

*Stegostoma tigrinum (Forster). Chiloscyllium indicum (Gmelin). Scoliodon walbeehmii Rhunchobatis djiddensis (Forskal). (Bleeker). *Mobula eregoodoo-tenke (Cuvier). Elops hawai-iensis Regan. Megalops cyprinoides (Broussonet). Chirocentrus dorab (Forskal). Chanos chanos (Forskal). Dussumieria hasseltii Bleeker. dinella fimbriata (Valenciennes). Ilisha hoevenii (Bleeker). Anodontostoma chacunda (B. Hamilton). Anchovia commersonii (Lacépède). socles hamiltoni (Gray). T. mystax (Schneider). T. setirostris (Broussonet). Saurida argyrophanes Richardson. *Monopterus albus (Zuiew). guilla mauritiana Bennett. Muraenesox cinereus (Forskal). Pisodonophis cancrivorus (Richardson). P. boro (B. Hamilton). *Evenchelys macrurus (Bleeker). Carassius auratus (Linnæus). Netuma thalassina (Rüppell). Clarias batrachus (Bloch). Tylosurus leiuroides (Bleeker). T. caudimaculatus (Čuvier). Hemiramphus limbatus Valenciennes. H. quoyi Valenciennes. *H. far (Forskal). H. cantoris Bleeker. Zenarchopterus dispar (Valenciennes). Z. philippinus (Peters). Dermatogenys viviparus (Peters). Liza labiosa (Valenciennes). L. waigiensis (Quoy and Gaimard). Sphyræna langsar Bleeker. Fistularia serrata Cuvier. Corythroichthys spicifer (Rüppell). Gastrotokeus biaculeatus (Bloch). Myripristis murdjan (Forskal). Holocentrus ruber Lacépède. *H. dia-

dema (Lacépède). H. cornutus Bleeker. Flammeo sammara (Forskal). Scomber microlepidotus Rüppell. Scomberomorus commerson Lacépède. Trichiurus haumela (Forskal). Scomberoides tolooparah (Rüppell). S. tala (Cuvier). Lactarius lactarius (Schneider). Megalaspis cordyla (Schneider). Decapterus kurra Bleeker. Selar boons (Cuvier). S. crumenophthalmus (Bloch). Caranx ignobilis (Forskal). C. sexfasciatus Quoy & Gaimard. C. kalla Cuvier. C. djeddaba (Forskal). C. hasseltii (Bleeker). C. leptolepis Cuvier. Carangoides oblongus (Valenciennes). Citula armata (Forskal). Scyris indica Rüppell. Leiognathus equula (Forskal). L. splendens (Cuvier). L. fasciatus (Lacépède). Deveximentum insidiator (Bloch). D. ruconius (B. Hamilton). Gazza minuta (Bloch). Amia quadrifasciata (Cuvier). A. kalloptera (Bleeker). A. fleurieu (Lacépède). A. koilomatadon (Bleeker). A. frenata (Valenciennes). *A. exostigma (Jordan & Seale). A. bifasciata (Rüppell). A. versicolor (Smith & Radcliffe). A. savayensis (Günther). A. jenkensi Evermann & Seale. *A. leptacantha (Bleeker). *Archamia zosterophora (Bleeker). *Cheilodipterus lineatus Lacépède. Ambassis ambassis (Lacépède). Priopus buruensis (Bleeker). Lates calcarifer (Bloch). Epinephelus bontoides (Bleeker). E. maculatus (Bloch). E. fasciatus (Forskal). Psammoperca waigiensis (Quov & Gaimard). *Pharopteryx corallicola (Rüppell). *Priacanthus tayenus Richardson. *Lutianus chirtah (Cuvier). L. decussatus (Cuvier). L. johni (Bloch). L. vitta (Quoy & Gaimard). L. lineolatus (Rüppell). Nemipterus japonicus (Bloch). N. luteus (Bloch). Pinjalo pinjalo (Bleeker). Caesio caerulaureus (Lacépède). Therapon theraps (Cuvier). T. puta (Cuvier). T. jarbua (Forskal). Helotes sexlineatus (Quoy & Gaimard). Mesopristes plumbeus (Kner). Pomadasis hasta (Bloch). P. maculatus (Bloch). Scolopsis

64 COPEIA

bilineatus (Bloch). Pentapus nemurus (Bleeker). P. caninus (Cuvier). *Plectorhinchus pictus (Thunberg). P. lineatus (Linnæus). Lethrinus harak (Forskal). L. moensi (Bleeker). L. leutjanus (Lacépède). L. mahsena (Forskal). Sparus berda (Forskal). Xystæma punctatum (Cuvier). X. oyena (Forskal). X. poieti (Cuvier). X. baconensis Evermann & Seale. X. kapas (Bleeker). *X. abbreviata (Bleeker). *X. macracantha (Bleeker). Argyrosomus aneus (Bloch). A. goldmani (Bleeker). Sciana dussumieri (Valenciennes). S. russelli (Cuvier). Sillago sihama (Forskal). Polydactylus plebius (Broussonet). Pseudochromis ta-peinosomus (Bleeker). Dampieria melanotaenia (Bleeker). *Pseudoplesiops typus Bleeker. Ophiocephalus striatus Bloch. Anabas testudineus (Bloch). Pomacentrus pavo (Bloch). P. violescens Bleeker. P. taeniurus Bleeker. P. lividus (Forster). *P. notophthalmus Bleeker. *Cheiloprion labiatus (Day). Abudefduf antjerius (Cuvier). A. coelestinus (Cuvier). A. saxatilis (Linnæus). Chromis caeruleus (Cuvier). Dascyllus aruanus (Linnæus). D. trimaculatus (Rüppell). Choerodon anchorago (Bloch). Stethojulis strigiventer (Bleeker). Halichoeres leparensis Bleeker. Cheilio inermis (Forskal). Hemipteronotus pentadactylus (Lacépède). Scatophagus argus (Gmelin). Drepane punctata (Gmelin). Platax orbicularis (Forskal). Chaetodon auriga Forskal. *C. fasciatus Forskal. C. punctato-fasciatus Valenciennes. C. trifasciatus Park. *Hemiochus varius (Cuvier). Zanclus cornutus (Linnæus). Hepatus celebicus (Bleeker). Naso unicornis (Forskal). Siganus fuscescens (Houttuyn). S. marmoratus (Quoy & Gaimard). S. vermiculatus (Quoy & Gaimard). S. javus (Linnæus). S. corallinus (Valenciennes). Triacanthus oxycephalus Bleeker. T. blochii Bleeker. Balistapus undulatus (Park). Monacanthus surothura Van Hasselt. Osbeckia COPEIA 65

scripta (Osbeck). Spheroides lunaris (Schneider). *Ostracion arcus Schneider. O. cornutum Lin-O. gibbosum Linnæus. Eleotris fusca (Schneider). Butis butis (B. Hamilton). Gobius ornatus Rüppell. Glossogobius giuris (B. Hamil-*Oxyurichthys tentacularis (Valenciennes). ton. *Gobiodon ceramensis Bleeker. Echeneis naucrates (Linnæus). Sebastopsis guamensis (Quoy & Gaimard). Pterois volitans (Linnæus). *Inimicus didactylum (Pallas). Platycephalus indicus (Linnæus). Thysanophrys crocodilus (Tilesius). T. bataviensis Bleeker. Elates thompsoni Jordan & Seale. Psettodes erumei (Bloch). Pseudorhombus javanicus (Bleeker). Cynoglossus sumatrensis (Bleeker). *C. quadrilineatus (Bleeker). Coryzichthys diemensis (Le Sueur). *Salarias fuscus Rüppell. *Petroscirtes hypseloptera Bleeker.

HENRY W. FOWLER, Philadelphia, Pa.

NOTE ON THE DUSKY SALAMANDER.

While spending a week-end at Slabsides, John Burroughs' rustic cabin near West Park, Ulster County, New York, I had an experience with the Dusky Salamander, Desmognathus fuscus (Raf.), which may be worth recording. After dark, on the night of November 25, 1917, I went to the spring to get a bucket of water, and when I returned to the light, I was surprised to find that I had dipped up two active immature salamanders. A few minutes later one of these was inadvertently poured out into a pan containing a little water, scarcely a half-inch deep, and allowed to remain in that all night. The next morning, the temperature having dropped to about 20° F., the salamander was frozen solid in the ice—that is, the ice was apparently frozen solid all around it. But after we had started a fire in the fireplace and the room had warmed up a bit, the ice thawed out, and the salamander, as soon as it was free from its icy prison, was as lively as it was the evening before, seemingly unharmed by the experience. In fact, after the two specimens were again placed together in water, it was impossible to tell which had been encased in the ice, as they were uniformly active and continued so during the several days that I kept them alive.

A single observation should not be taken to indicate too much, but this one points to nocturnal, or at least crepuscular